Ref 71c

INTERNATIONAL INSTITUTE OF AGRICULTURE

INTERNATIONAL BULLETIN OF PLANT PROTECTION

PUBLISHED MONTHLY

REPRINT FROM THE 'INTERNATIONAL REVIEW OF AGRICULTURE'

YEAR XIX – 1945

ROME
VILLA UMBERTO I
1945

STAFF OF THE PLANT PROTECTION SECTION

Chief of Section: Professor Dr GIULIO TRINCHIERI

Councillor: Dr NORBERT VON GESCHER

Page numbers of the different issues.

January & February:
(Nos. 1 & 2) pages 1 M to 16 M.

March & April:
(Nos. 3 & 4) pages 17 M to 32 M.

May & June:
(Nos. 5 & 6) pages 33 M to 48 M.

July & August:
(Nos. 7 & 8) pages 49 M to 72 M.

September & October):
(Nos. 9 & 10) pages 73 M to 88 M.

November & December:
(Nos. 11 & 12) pages 89 M to 112 M.

INDEX TO 'THE INTERNATIONAL BULLETIN OF PLANT PROTECTION' FOR THE YEAR 1945

I. — Index of Authors.

BALDACCI, E., see CIFERRI, R. and BALDACCI, E.

CAVALLI, L., and CIFERRI, R. Results of the statistical analysis of six distributions of germination percentages of conidia of *Alternaria, tenuis* type 92 M.

CIFERRI, R. Checking of accuracy in testing fungicides 'in vitro', and examples particularly as applied to *Alternaria, tenuis* type, 17 M.

— and BALDACCI, E. Comparative experiments on the toxicity of different metals on *Alternaria, tenuis* type, and *Plasmopara viticola*, 33 M.

— See also CAVALLI, L., and CIFERRI, R.

JANNONE, G. Serious decay of papaw caused by a nema, in Eritrea, 76 M.

— First observations on a mite parasite of the papaw and on a predator of same, in Eritrea, 90 M.

MINISTRY OF AGRICULTURE AND FORESTS, ROME. Colorado beetle (*Leptinotarsa decemlineata*) infestation in Piedmont, Italy, 89 M.

SILVESTRI, F. Contribution to the biology of the olive gall gnat (*Prolasioptera berlesiana* Paoli) in Italy, 73 M.

TRINCHIERI, G., Air transport and phytosanitary control, 1 M.

— Third contribution to the locust bibliography, 22 M, 38 M, 100 M.

— A copy of the *Sylloge fungorum* with corrections, additions, annotations and other details appended by the author, 49 M.

— Material for an European Colorado beetle bibliography, 81 M, 109 M.

II. — Subject Index.

Acrospermum conicum, 59 M.

Acrothecium Tulasnei, 58 M.

Acurtis, 63 M.

Aecidium strobilinum, 61 M, 63 M.

Agromyza, 5 M.

Agrotis ypsilon, 8 M.

Aircraft Pyagra (insecticide), 12 M.

Air transport and phytosanitary control, 1 M.

Alabama argillacea, 8 M.

Aleurites triloba : Diplodia, 6 M.

Aleurocanthus woglumi (Citrus whitefly), 4 M, 5 M, 6 M.

Alternaria brassicae, see *Alternaria herculea*.

— *herculea* (*A. brassicae*), 6 M.

— *tenuis* type, 17 M, 19 M, 20 M, 33 M, 57 M, 92 M.

Amblyosporium Botrytis, 65 M.

American mildew of gooseberry, see *Sphaerotheca mors-uvae*.

Amorbia emigratella, 6 M.

Amphisphaeria zerbina, 57 M.

Ananas sativus (pineapple) : *Leucopholis irrorata*, 7 M.

Anastrepha, 6 M.

— *ludens*, 6 M.

Antennaria scoriadea, 53 M.

Anthostomella Smilacis, 54 M.

Anthracnose fungus of citrus, see *Colletotrichum gloeosporioides*.

Antirrhinum : Frankliniella intonsa, 5 M.

— — — f. *adusta*, 5 M.

— Geometrid, 5 M.

— *Haplothrips aculeatus*, 5 M.

— Mirid, 6 M.

— *Phytometra* (*Plusia*) *gamma*, 5 M.

Antirrhinum : Pterophorid, 6 M.
— Syrphid, 6 M.
— *Taeniothrips atratus*, 5 M.
— *Thrips flavus*, 5 M.
— — *fuscipennis*, 5 M.
— — — f. *corticina*, 5 M.
— — *validus*, 5 M.
Anychus latus on papaw in Eritrea, 90 M.
Arachis glabrata : *Phyllosticta*, 6 M.
Argentine ant, see *Iridomyrmex humilis*
Arrhenia (*Apus*) *fimicola*, 63 M.
— *tenella*, 69 M.
Arthopyrenia Aspiciliae, 55 M.
Aschersonia, 60 M.
Ascochyta arenaria, 67 M.
Asinapta furcifer, 76 M.
Asparagus beetle, see *Crioceris asparagi*.
Asparagus : *Crioceris asparagi* (asparagus beetle), 3 M.
Aspidiotus palmae, 6 M.
— *persearum*, 6 M.
— *spinosus*, 6 M.
Assyrian plum, see *Cordia myxa*.
Aster : *Coleophora*, 5 M.
— *Frankliniella intonsa*, 5 M.
— *Phytometra* (*Plusia*) gamma, 5 M.
— Syrphid, 6 M.
— *Taeniothrips atratus*, 5 M.
— *Thrips fuscipennis*, 5 M.
— *Tritogenaphis* (*Macrosiphum*), 5 M.
Asterina, 6 M.
Asteroma Dentariae, 67 M.
— *radiatum*, 67 M.
Belgium : Legislative and administrative measures, 78 M (protection of the production of tested seed and seedlings) ; 79 M (fungicides, insecticides, weedicides and parasiticides ; Colorado beetle, *Leptinotarsa decemlineata*) ; 80 M (capture of sparrows).
Bibliography :
Colorado beetle, 81 M, 109 M.
Locusts, 22 M, 38 M, 100 M.
Black rot of the grape vine, see *Guignardia bidwellii*.
Blissus leucopterus (chinch bug), 8 M.
Borrichia glabrata : *Frankliniella insularis*, 6 M.
Botryosporium pulchrum, 61 M.
Botrytis umbellata, 61 M.
Brassica oleracea : *Alternaria herculea* (*A. brassicae*), 6 M.
Cabbage : *Pieris rapae* (small cabbage butterfly), 3 M.
Calonectria Otagensis, 58 M.
Camellia : *Aspidiotus palmae*, 6 M.
— — *persearum*, 6 M.
— *Aspidiotus spinosus*, 6 M.
— *Chrysomphalus personatus*, 7 M.
— *japonica* : *Capnodium*, 6 M.
— — *Cephaleuros virescens*, 6 M.
Canavalia obtusifolia : *Frankliniella insularis*, 6 M
Canna : *Leucopholis irrorata*, 7 M.
Cantharellus aurantiacus, 69 M.
Capnodium, 6 M.
— *Footii*, 53 M.
Carica papaya (papaw) : *Anychus latus* and *Scymnus morelleti* in Eritrea, 90 M.
— — — *Heterodera marioni* (*H. radicicola*) in Eritrea, 76 M.
Carnation : *Heterosporium echinulatum* (carnation ring spot), 5 M.
— ring spot, see *Heterosporium echinulatum*.
Cassia siamea (Siamese senna) : *Leucopholis irrorata*, 7 M.
Cephaleuros virescens, 6 M.
Ceratium hydnoides, 69 M.
Cercospora, 6 M.
Ceriospora zantha, 57 M.
Chinch bug, see *Blissus leucopterus*.
Chrysanthemum : *Septoria chrysanthemi* (*S. chrysanthemella*) (chrysanthemum blotch), 5 M.
Chrysanthemum blotch, see *Septoria chrysanthemi* (*S. chrysanthemella*).
Chrysomphalus personatus, 6 M.
Circinotrichum murinum, 62 M.
Citrullus vulgaris (water melon) : *Leucopholis irrorata*, 7 M.
Citrus : *Aleurocanthus woglumi* (citrus white fly), 5 M.
— *Coccus viridis* (green coffee scale), 6 M.
— *Diaporthe citri*, 6 M.
— *Phomopsis citri* (citrus melanose), 6 M.
— *aurantium* : *Phomopsis citri*, 6 M.

Citrus aurantium : Sphaceloma fawcettii, 6 M.
— *grandis* : *Lepidosaphes beckii*, 6 M.
— — *Pseudaonidia* (*Selenaspidus*) *articulatus*, 6 M.
— *nobilis* var. *deliciosa* : *Sphaceloma fawcettii* (citrus scale), 6 M.
— *sinensis* : *Anastrepha ludens* (Mexican fruit fly), 6 M.
— — *Colletotrichum gloeosporioides* (anthracnose fungus of citrus), 6 M.
— — *Lepidosaphes beckii* (citrus mussel scale), 6 M.
— — *Septobasidium*, 6 M.
Citrus melanose, see *Phomopsis citri*.
— mussel scale, see *Lepidosaphes beckii*.
— scab, see *Sphaceloma fawcettii*.
— white-fly, see *Aleurocanthus woglumi*.
Clathrosphaera spirifera, 61 M.
Clavaria minuta, 69 M.
— *palmata*, 69 M.
Coccus viridis (green coffee scale), 6 M, 7 M.
Coffea : *Aleurocanthus woglumi* (citrus white fly), 6 M.
— *Coccus viridis* (green coffee scale), 7 M.
Coleophora, 5 M.
Colletomanginia, 71 M.
Colletotrichum, 6 M.
— *gloeosporioides* (anthracnose fungus of citrus), 6 M.
Colorado beetle, see *Leptinotarsa decemlineata*.
Coniothyrella agaricola, 72 M.
Coniothyrina agaricola, 72 M.
Coniothyrium Cargillianum, 70 M.
Cordia myxa (Assyrian plum) : *Leucopholis irrorata*, 8 M.
Corticium amorphum, 69 M.
Cortinarius elatior, 69 M.
— *Friesii*, 63 M.
— *lilacinus*, 69 M.
— *montellicus*, 69 M.
Corynelia uberata, 62 M.
Cotton bollworm, 9 M.
— *Pectinophora* (*Platyedra*) *gossypiella* (pink cotton bollworm), 6 M.
Crioceris asparagi (asparagus beetle), 3 M.
Cronartium ribicola (white pine blister rust), 3 M.
Cucurbitaria Rutae, 58 M.
— *solitaria*, 58 M.
Cytospora stenospora, 59 M.
Dacus oleae (olive fly), 73 M.
Dahlia : *Frankliniella intonsa*, 5 M.
— Geometrid, 5 M.
— *Thrips fuscipennis*, 5 M.
Darluca Filum, 59 M.
Dasyscypha calycina, 69 M.
— *willkommii* (European larch canker), 3 M.
Dematium articulatum, 62 M.
— *hispidulum*, 62 M.
Deo-Base (insecticide), 13 M.
Deskito (insecticide), 12 M.
Dianthus : *Cercospora*, 6 M.
— *Heterosporium echinulatum* (*Didymellina dianthi*), 6 M.
— Mirid, 5 M.
— *Saissetia hemisphaerica*, 5 M.
— *Taeniothrips atratus*, 5 M.
— *Thrips fuscipennis* f. *corticina*, 5 M.
— — *validus*, 5 M.
Diaporthe citri, 6 M.
— *Chorostate fibrosa*, 55 M.
— — *oxyspora*, 55 M.
— — *perjuncta*, 55 M.
— — *strumellaeformis*, 55 M.
— — *Tessella*, 55 M.
— *Euporthe ceuthosporioides*, 55 M.
— — *discors*, 55 M.
— — *Tulasnei*, 55 M.
— — *viticola*, 55 M.
— *Tetrastaga striaeformis*, 56 M.
Diatrype platystoma, 54 M.
Diatrypella, 54 M.
Dichaena quercina, 59 M.
Didymella glomerulata, 55 M.
Didymellina dianthi, see *Heterosporium echinulatum*.
Didymium hypnophilum, 69 M.
— *squamulosum*, 69 M.
Didymosphaeria (*Microthelia*), 70 M.
— *parietinaria*, 70 M.
— *Puyae*, 70 M.
— *rugulosaria*, 70 M.

Didymosphaeria stereocaulicola, 70 M.
— *stictaria*, 70 M.
Dimerosporium abjectum, 53 M.
— *Ellisii*, 53 M.
Dinemasporium graminum, 60 M.
— *strigosum*, 60 M.
Diospyros discolor (mabolo) : *Leucopholis irrorata*, 7 M.
Diplocarpon rosae (rose black spot), 6 M.
Diplodia, 6 M.
Downy mildew of the vine, see *Plasmopora viticola*.
Drosophila melanogaster (grape fruit fly), 8 M.
— *repleta*, 8 M.
Dwarfing, arice virus disease caused by *Nephotettix*, 7 M.
Elasmopalpus lignosellus (lesser corn stalk borer), 8 M.
Elm : *Ophiostoma ulmi* (elm disease), 3 M.
Elm disease, see *Ophiostoma ulmi*.
Epicymatia, 65 M.
— *lichenicola*, 55 M.
— *vulgaris*, 55 M.
Epidendrum : *Colletotrichum*, 6 M.
Eritrea : *Anychus latus* dangerous to papaw, 90 M.
— *Heterodera marioni* injurious to papaw, 76 M.
Erysiphe communis, 52 M.
— *lamprocarpa*, 52 M.
— *Martii*, 52 M.
Eugenia jambolana (jambolan plum) : *Leucopholis irrorata*, 7 M.
European corn borer, see *Pyrausta nubilalis*.
— larch canker, see *Dasyscypha willkommii*.
Eurotium stercorarium, 53 M.
Euryachora stellaris, 58 M.
Eutypa bellula, 54 M.
— *lejoplaca*, 54 M.
— *phaselina*, 54 M.
— *subtecta*, 54 M.
Euxesta notata (spotted root fly), 8 M.
Excipula strigosa, 60 M.
— — *hysteriiformis*, 60 M.
Exidiopsis, 66 M.
— *effusa*, 66 M.
Exogone Kaiseriana, 72 M.
Fall armyworm, see *Laphygma frugiperda*.
Frankliniella cephalica, 7 M.
— *cubensis*, 7 M.
— *fortissima*, 7 M.
— *insularis*, 6 M.
— *intonsa*, 5 M.
— — f. *adusta*, 5 M.
Fumago, 53 M.
Gaillardia grandiflora : *Frankliniella intonsa*, 5 M.
— — *Taeniothrips atratus*, 5 M.
— — *Thrips fuscipennis*, 5 M.
Galactinia proteana var. *sparassoides*, 72 M.
Gardenia : *Coccus viridis* (green coffee scale), 6 M.
Gardenia florida : *Amorbia emigratella*, 6 M.
— — *Coccus viridis* (green coffee scale), 6 M.
— — *Macrosporium*, 6 M.
— — *Meliola*, 6 M.
— — *Protopulvinaria* (*Pulvinaria pyriformis*), 6 M.
— — *Saissetia oleae* (olive scale), 6 M.
— — *Septobasidium*, 6 M.
— — *Stilbum*, 6 M.
Geometrid, 5 M.
Gibberella polycocca, 58 M.
Gladiolus : *Frankliniella fortissima*, 7 M.
— Mirid, 5 M.
— *Taeniothrips atratus*, 5 M.
— — *simplex*, 7 M.
Gloeosporium Betulae, 61 M.
— *Salicis*, 61 M.
Glomerella, 67 M.
Gnomoniopsis, 67 M.
Gooseberry : *Sphaerotheca mors-uvae* (American mildew of gooseberry), 3 M.
Grapefruit fly, see *Drosophila melanogaster*.
Grape vine : *Guignardia bidwellii* (black rot of the grape vine), 3 M.
Green coffee scale, see *Coccus viridis*.
— peach aphis, see *Myzus persicae*.
Grey wood of Seroat, see *Streblus asper*.
Guignardia bidwellii (black rot of the grape vine), 3 M.

Guinea grass, see *Panicum maximum*.
Gymnosporangium gracile, 68 M.
Haplothrips aculeatus, 5 M.
Hebeloma elatum, 69 M.
Helicomyces tubulosus, 61 M.
Helicotrichum obscurum, 62 M.
Heliothis obsoleta, 8 M.
Hendersonia Sabaleos, 60 M.
— *uredinicola*, 59 M.
Heterodera marioni, 77 M.
— *radicicola*, 77 M.
Heterosporium echinulatum (carnation ring spot), 5 M.
Hippotion celerio, 8 M.
Hymenula, 62 M.
Hyphaene nodularia, 91 M.
Hypocreopsis riccioidea, 58 M.
Illosporium roseum, 62 M.
Indian meal moth, see *Plodia interpunctella*.
— pod borer, see *Maruca testulalis*.
Inzengaea, 64 M.
— *erythospora*, 65 M.
Iridomyrmex humilis (Argentine ant), 2 M, 3 M.
Italy : Legislative and administrative measures, 80 M (Royal Observatory for plant diseases for Calabria).
— Olive gall gnat (*Prolasioptera berlesiana*), 73 M.
— Colorado beetle (*Leptinotarsa decemlineata*), 89 M.
Jambolan plum, see *Eugenia jambolana*.
Japanese beetle, see *Popillia japonica*.
Jasminum sambac : Phomopsis, 6 M.
— — *Taeniothrips hawaiiensis*, 7 M.
Lachnocladium Mollerianum, 64 M.
Lactarius aurantiacus, 69 M.
— *pallidus*, 69 M.
Laestadia, 54 M.
Laphygma frugiperda (fall armyworm), 8 M.
Larmuth ejector, 12 M.
Lecanium corni (plum scale), 5 M.
Legislative and administrative measures :—
Belgium, 78 M, 79 M, 80 M.
Italy, 80 M.
Lema melanopa, 8 M.
Lentinus auricula, 69 M.
Lentinus badius, 69 M.
— *bissus*, 69 M.
Lepidosaphes beckii (citrus mussel scale), 6 M.
Lepiota candida, 71 M.
— *pulcherrima*, 71 M.
Leptinotarsa decemlineata (Colorado beetle), 2 M, 3 M, 81 M, 89 M, 109 M.
Leptosphaeria baeomycearia, 70 M.
— *circinans*, 57 M.
— *Sabauda*, 57 M.
— *Umbilicariae*, 70 M.
Leptothyrium Betulae, 61 M.
Lesser corn stalk borer, see *Elasmopalpus lignosellus*.
Lettuce : *Tritogenaphis* (*Macrosiphum*) *sonchi* (lettuce aphis), 5 M.
— aphis, see *Tritogenaphis* (*Macrosiphum*) *sonchi*.
Leucopholis irrorata, 7 M.
Lichenopsis sphaeroboloidea, 60 M.
Limothrips denticornis, 7 M.
Locusts : Third contribution to the l. bibliography, 22 M, 38 M, 100 M.
Lychnis : Agromyza, 5 M.
— *Frankliniella intonsa*, 5 M.
— *Taeniothrips atratus*, 5 M.
— *Thrips fuscipennis*, 5 M.
Lygaeus pandurus, 8 M.
Macrophoma leucostigma, 66 M.
Macrosiphum luteum, 7 M.
— See also *Tritogenaphis* (*Macrosiphum*) *sonchi*.
Macrosporium commune, 57 M.
Maize, see *Zea mays*.
Malbranchea, 61 M.
Mammea : Aspidiotus spinosus, 6 M.
Mangifera indica (mango) : *Leucopholis irrorata*, 7 M.
Mango, see *Mangifera indica*.
Mangoes : *Anastrepha*, 6 M.
Maruca testulalis (Indian pod borer), 8 M.
Massaria, 57 M.
Mazzantia circumscripta, 58 M.
Melanomma dubiosum, 57 M.
— *lonicerinum*, 57 M.
Melanospora rhizophila, 71 M.
Meliola quinquespora, 53 M.
Mentha : Coccus viridis (green coffee scale), 6 M.

Metals : Toxicity of different m. on *Alternaria, tenuis* type, and *Plasmopara viticola*, 33 M.
Metasphaeria nephromiaria, 70 M.
Mexican fruit fly, see *Anastrepha ludens*.
Microsphaera Platani, 52 M.
— *Viburni*, 52 M.
Microthelia atricola, 70 M.
— *baeomycearia*, 70 M.
— *cargilliana*, 70 M.
— *collemaria*, 70 M.
— *Cookei*, 70 M.
— *nephromiaria*, 70 .
— *parietinaria*, 70 M.
— *perrugosaria*, 70 M.
— *Puyae*, 70 M.
— *ramalinaria*, 70 M.
— *rugulosaria*, 70 M.
— *stereocaulicola*, 70 M.
— *stictaria*, 70 M.
— *Umbilicariae*, 70 M.
— *vesicularia*, 70 M.
Microthyrium epimyces, 65 M, 68 M.
Mirid, 5 M.
Monilia albolutea, 65 M.
Mycena speirea, 62 M.
Mycosphaerella Cookei, 70 M.
Myzus persicae (green peach aphis), 5 M.
Nectria armeniaca, 61 M.
Nephotettix (rice leafhopper), 7 M.
Nitschkea Flageoletiana, 68 M.
Olive fly, see *Dacus oleae*.
Olive gall gnat, see *Prolasioptera berlesiana*.
Olive scale, see *Saissetia oleae*.
Onion thrips, see *Thrips tabaci*.
Orchids : *Asterina*, 6 M.
— *Macrosiphum luteum*, 7 M.
— *Phoma*, 6 M.
Ophiostoma ulmi (elm disease), 3 M.
Oryza sativa (rice) : *Leucopholis irrorata* and *Nephotettix*, 7 M.
Pachymerus pallidus, 8 M.
Panicum maximum (Guinea grass) : *Leucopholis irrorata*, 7 M.
Papaw, see *Carica papaya*.
Peach : *Sphaerotheca pannosa* (peach mildew), 6 M.
Peas, see *Pisum sativum*.
Pectinophora (*Platyedra*) *gossypiella* (pink cotton bollworm), 6 M.
Peziza aurantia, 64 M.
— (*Elvela*) *coccinea*, 64 M.
— (*Sarcoscypha*) *coccinea*, 64 M.
Phantomyst, 12 M.
Pharcidia, 65 M.
Phoma, 6 M.
— *Berberidis*, 59 M, 65 M.
— *Brassicae*, 59 M.
— *dalmatica*, 73 M.
— *herbarum*, 57 M.
— *leucostigma*, 66 M.
— *Lingam*, 59 M.
— *Roumeguerii*, 59 M.
Phomopsis citri (citrus melanose), 6 M.
Phthorimaea operculella (potato tuber moth), 2 M.
Phyllachora Campanulae, 58 M.
Phyllosticta, 6 M.
Phylloxera of the vine, see *Phylloxera vitifoliae*.
Phylloxera vitifoliae (phylloxera of the vine), 3 M.
Phymatosphaera abyssinica, 53 M.
Phytometra (*Plusia*) *gamma*, 5 M.
Phytophthora infestans (potato blight), 3 M.
Phytosanitary control and air transport, 1 M.
Pieris rapae (small cabbage butterfly), 3 M.
Piggotia astroidea, 60 M.
Pineapple, see *Ananas sativus*.
Pink cotton bollworm, see *Pectinophora* (*Platyedra*) *gossypiella*.
Pisum sativum (peas) : *Leucopholis irrorata*, 7 M.
Placosphaeria, 59 M.
Plasmopara viticola (downy mildew of the vine), 3 M, 33 M.
Platyedra gossypiella (pink cotton bollworm), 9 M.
Platyedra, see *Pectinophora* (*Platyedra*) *gossypiella*.
Pleospora herbarum, 57 M.
— *infectoria*, 57 M.
— *vulgaris*, 57 M.
Pleosporopsis strobilina, 61 M.

Pleosporopsis strobilorum, 61 M.
Plodia interpunctella (Indian meal moth), 8 M.
Plum : *Lecanium corni* (plum scale), 5 M.
Plumeria acutifolia : *Taeniothrips hawaiiensis*, 7 M.
Plusia, see *Phytometra* (*Plusia*) *gamma*, 5 M.
Plutella maculipennis, 8 M.
Podagrica puncticollis, 8 M.
Polyporus, 66 M.
— *niveus*, 63 M.
Popillia japonica (Japanese beetle), 9 M, 13 M.
Potato : *Leptinotarsa decemlineata* (Colorado beetle), 2 M, 81 M, 109 M.
— *Phthorimaea operculella* (potato tuber moth), 2 M.
— *Phytophthora infestans* (potato blight), 3 M.
— *Synchytrium endobioticum* (wart disease), 3 M.
Potato blight, see *Phytophthora infestans*.
— tuber moth, see *Phthorimaea operculella*.
Powdery mildew of the vine, see *Uncinula necator*.
Prolasioptera berlesiana (olive gall gnat), 73 M.
Prosthemiella formosa, 61 M.
Protomyces pachydermus, 63 M.
Protopulvinaria (*Pulvinaria*) *pyriformis*, 6 M.
Pseudaonidia (*Selenaspidus*) *articulatus* (rufous scale), 6 M.
Psilopezia, 68 M.
Pterophorid, 6 M.
Pulvinaria, see *Protopulvinaria* (*Pulvinaria*) *pyriformis*.
Puncture vine, see *Tribulus terrestris*.
Purple mombin, see *Spondias purpurea*.
Pyrausta nubilalis (European corn borer), 3 M.
Pyrenochaeta Berberidis, 59 M, 65 M.
Pyrenopeziza Campanulae, 58 M.
— *Phyteumatis*, 58 M.
Pyrenophora relicina, 58 M.
Rice (*Oryza sativa*) : *Tylenchus angustus* (*Anguillulina angusta*), causal agent of the rice disease called Ufra, 5 M.
— leafhopper, see *Nephotettix*.
— *Leucopholis irrorata* 7 M.
Rosa (rose) : *Chrysomphalus personatus*, 7 M.
— *Diplocarpon rosae* (rose black spot), 6 M.
— *Franklíniella cephalica*, 7 M.
— — *cubensis*, 7 M.
— *Limothrips denticornis*, 7 M.
— *Sphaceloma rosarum* (rose leaf spot), 5 M.
— *Sphaerotheca pannosa* (rose mildew), 6 M.
— *Taeniothrips ericae*, 7 M.
— *Typhlocyba rosae* (rose leafhopper), 5 M.
Rose black spot, see *Diplocarpon rosae*.
— leafhopper, see *Typhlocyba rosae*.
— leaf spot, see *Sphaceloma rosarum*.
— mildew, see *Sphaerotheca pannosa*.
Rosellinia, 61 M.
— *sepulta*, 64 M.
— *Tassiana*, 54 M.
Rufous scale, see *Pseudaonidia* (*Selenaspidus*) *articulatus*.
Ruhlandiella, 67 M, 68 M.
Saccardia Martini, 53 M.
Saccharum officinarum (sugarcane) : *Leucopholis irrorata*, 7 M.
Saissetia hemisphaerica, 5 M.
— *oleae* (olive scale), 6 M.
Sandal, see *Sandoricum indicum*.
Sandoricum indicum (sandal) : *Leucopholis irrorata*, 7 M.
Scleroderma verrucosum, 63 M.
— *vulgare*, 63 M.
Sclerotinia mycetophila, 65 M.
Scymnus morelleti, a predator of *Anychus latus* on papaw in Eritrea, 91 M.
Selenaspidus, see *Pseudaonidia* (*Selenaspidus*) *articulatus*.
Selenothrips rubrocinctus, 7 M.
Sepedonium agglomeriferum, 61 M.
Septobasidium, 6 M.
Septoria Calystegiae, 60 M.
— *Caraganae*, 59 M.

Septoria chrysanthemi (*S. chrysanthemella*) (chrysanthemum blotch), 5 M.
— *Leucanthemi*, 60 M.
— *pallens*, 60 M.
— *Theae*, 65 M.
— *Veronicae*, 60 M.
Siamese senna, see *Cassia siamea*.
Sinoxylon ruficorne, 8 M.
Small cabbage butterfly, see *Pieris rapae*.
Sphaceloma fawcettii (citrus scale), 6 M.
— *rosarum* (rose leaf spot), 5 M.
Sphaerella caulicola, 71 M.
— *Ceratoniae*, 55 M.
— *Hermione*, 55 M.
— *Leguminis-Cytisi*, 55 M.
— *Pseudacaciae*, 55 M.
— *subcongregata*, 65 M.
Sphaeria decorticans, 55 M.
— *fusispora*, 57 M.
— *lichenicola*, 55 M.
— *Solani*, 56 M.
— *versatilis*, 55 M.
Sphaeropsis Dearnessii, 72 M.
— *ontariana*, 72 M.
Sphaerosoma, 68 M.
Sphaerotheca mors-uvae (American mildew of gooseberry), 3 M.
— *Niesslii*, 52 M.
— *pannosa* (peach and rose mildew), 6 M.
Sphaerulina intermixta, 55 M.
Spondias purpurea (purple mombin) : *Leucopholis irrorata*, 7 M.
Spotted root fly, see *Euxesta notata*.
Stictocephala festina (three-cornered alfalfa hopper), 8 M.
Streblus asper (grey wood of Seroat) : *Leucopholis irrorata*, 8 M.
Sugarcane, see *Saccharum officinarum*.
Sylloge fungorum omnium hucusque cognitorum : a copy of this work with corrections, additions, annotations and other details appended by the author, 49 M.
Syncladium, 53 M.
Synchytrium endobioticum (wart disease of the potato), 3 M.
Syrphid, 6 M.
Taeniothrips atratus, 5 M, 7 M.
Taeniothrips ericae, 7 M.
— *hawaiiensis*, 7 n.
— *simplex* (gladiolus thrips), 7 M.
Tamarind, see *Tamarindus indica*.
Tamarindus indica (tamarind) : *Leucopholis irrorata*, 8 M.
Teichospora princeps, 57 M.
Tetranychus telarius, 91 M.
Tetrastichus invidus, 76 M.
Thelephora, 69 M.
Three-cornered alfalfa hopper, see *Stictocephala festina*.
Thrips flavus, 5 M.
— *fuscipennis*, 5 M.
— — f. *corticina*, 5 M.
— *tabaci* (tobacco thrips), 5 M.
— *validus*, 5 M.
Tichothecium collemarium, 70 M.
— *erraticum*, 70 M.
— *pygmaeum*, 70 M.
Tobacco : *Thrips tabaci* (tobacco thrips), 5 M.
Trabutia quercina, 55 M.
Trematophaeria Alaterni, 57 M.
— *irregularis*, 57 M.
— *princeps*, 57 M.
Tribulus terrestris (puncture vine), 7 M.
Tripospora Cookei, 62 M.
Tritogenaphis (*Macrosiphum*) *sonchi* (lettuce aphis), 5 M.
Tylostoma Mollerianum, 64 M.
Typhlocyba rosae (rose leafhopper), 5 M.
Ufra, a rice disease caused by *Tylenchus angustus* (*Anguillulina angusta*), 5 M.
Uncinula columbiana, 66 M.
— *necator* (powdery mildew of the vine), 3 M.
Underwoodia Campbelli, 72 M.
Uredo fissa, 64 M.
— *fructicola*, 64 M.
— *pinguis*, 64 M.
Uromyces appendiculatus, 64 M.
Valsa diatrypa, 53 M.
Valsaria decorticans, 55 M.
Valsella clypeata, 53 M.
Vine : *Phylloxera vitifoliae* (phylloxera of the vine), 3 M.
— *Plasmopara viticola* (downy mildew of the vine), 3 M.

Wart disease of the potato, see *Synchytrium endobioticum*.
Water melon, see *Citrullus vulgaris*.
White pine blister rust, see *Cronartium ribicola*.
Xylaria contracta, 54 M.
Xylaria Poiteana, 54 M.
— *vernicosa*, 64 M.
Zasmidium scoriadeum, 53 M.
Zea mays (maize) : *Leucopholis irrorata*, 7 M.
Zingiber : Taeniothrips hawaiiensis, 7 M.

Prof. UGO PAPI, *General Secretary of the International Institute of Agriculture*

Printed by F. FAILLI, Printing-Office of the International Institute of Agriculture
ROME, Villa Umberto I, December 31, 1945.